



- 1a) Find the coordinates of the point lying between A (2,3) and B (8, -3) which divides the line segment AB in the ratio 1:2.

$$\begin{array}{l} 2 \rightarrow 8 \quad +6 \quad \frac{1}{3} = 2 \quad 2+2 = 4 \\ 3 \rightarrow -3 \quad -6 \quad \frac{1}{3} = -2 \quad 3-2 = 1 \end{array} \quad \underline{\underline{(4,1)}}$$


- b) Find the x-coordinate of the point where the perpendicular bisector of the line segment joining the points (2,-6) and (5,4) cuts the x-axis.



- c) The perpendicular bisector of the line segment joining the points (3,-5) and (1,1) passes through the point with coordinates (11,a). Find the value of a.




- d) The straight line  $L_1$  passes through points A (13,5) and B (9,2) and D.  
 The straight line  $L_2$  passes through points C (2,3) and D and is perpendicular to  $L_1$   
 Find the coordinates of D.




- e) Find the shortest distance between the parallel lines with equations:


$$x + 2y = 10 \quad \text{and} \quad x + 2y = 20$$



- f) A line  $L$  has equation  $y = 4 - 3x$ . A second line is perpendicular to  $L$  and passes through  $(-2, 0)$ . Find the area of the region enclosed by the two lines and the  $x$ -axis.




- g) The points  $A$ ,  $B$  and  $C$  have coordinates  $(0, 3)$  and  $(2, -1)$  and  $(k, 1)$  respectively.  $AB$  and  $BC$  are perpendicular. Find the area of the triangle  $ABC$ .




- h) The straight line  $L$  passes through points  $(2, 5)$  and  $(-2, 3)$  and meets the coordinate axes at  $P$  and  $Q$ . Find the area of a square with side  $PQ$ .

gradient  $L = \frac{2}{4} = \frac{1}{2}$   $|PQ|^2 = 4^2 + 8^2$   
 $y - 5 = \frac{1}{2}(x - 2)$   $= 16 + 64$   
 $y = \frac{1}{2}x + 4$   $= 80$   
 $P(0, 4)$   $Q(-8, 0)$


- i) The points  $A$  and  $B$  have coordinates  $(-1, 4)$  and  $(3, -2)$  respectively. A line  $L$  is perpendicular to  $AB$  and passes through  $B$ . Find the area of the region enclosed by  $L$  and the coordinate axes.



- j) The points  $A$  and  $B$  have coordinates  $(-4,5)$  and  $(0,4)$  respectively.  
The point  $C$  lies on the straight line through  $A$  and  $B$  such that the distance  $AB$  is the same as the distance  $BC$ . Find the coordinates of  $C$ .



- k) The points  $A$  and  $B$  have coordinates  $(1, 4\sqrt{3})$  and  $(-3 + \sqrt{3}, 3)$  respectively.  
The straight line  $L$  through  $A$  and  $B$  meets the  $x$ -axis at  $C$ .  
Calculate the acute angle between  $L$  and the  $x$ -axis




$$\tan \theta = \frac{4\sqrt{3} - 3}{1 + 3 - \sqrt{3}} = \frac{4\sqrt{3} - 3}{4 - \sqrt{3}}$$

$$= \frac{\sqrt{3}(4 - \sqrt{3})}{4 - \sqrt{3}} = \sqrt{3}$$

$$\theta = 60^\circ$$

- l) The points  $A$  and  $B$  have coordinates  $(8,2)$  and  $(11,3)$  respectively.  
The point  $C$  lies on the straight line with equation  $x + y = 14$   
Given that the distance  $AC$  is twice as large as the distance  $AB$ , find the two possible sets of coordinates of  $C$ .



$$|AB| = \sqrt{1^2 + 3^2} = \sqrt{10}$$

$$|AC| = 2\sqrt{10} \quad AC^2 = 40$$

$$40 = (a - 8)^2 + (12 - a)^2$$

$$40 = a^2 - 16a + 64 + 144 - 24a + a^2$$

$$2a^2 - 40a + 168 = 0$$

$$a^2 - 20a + 84 = 0$$

$$(a - 6)(a - 14) = 0$$

$$a = 6, 14$$


$C(a, 14 - a)$   
 $(6, 8)$   
 $(14, 0)$

- 2a) The straight line segment joining the points (6,-3) and (14,9) is a diameter of a circle. What is the equation of the circle?

$$\begin{aligned} \text{Midpoint} &= (10, 3) \\ \text{Diameter} &= \sqrt{12^2 + 8^2} \\ &= \sqrt{144 + 64} \\ &= \sqrt{208} \\ &= 2\sqrt{52} \end{aligned} \quad \begin{aligned} \text{Radius} &= \sqrt{52} \\ r^2 &= 52 \\ (x-10)^2 + (y-3)^2 &= 52 \end{aligned}$$

- b) The straight line segment joining the points <sup>A</sup>(-4,3) and <sup>B</sup>(0,5) is a chord of a circle with centre on the line with equation  $y = 3x + 5$ . What is the equation of the circle?


$$\begin{aligned} \text{gradient } AB &= \frac{1}{2} & \text{gradient } L &= -2 \\ \text{midpoint } AB &= (-2, 4) & y - 4 &= -2(x + 2) \\ & & y &= -2x \\ -2x &= 3x + 5 & r^2 &= 3^2 + 1^2 = 10 \\ -5 &= 5x & (x+1)^2 + (y-2)^2 &= 10 \\ x &= -1, y = 2 & \text{centre } &(-1, 2) \end{aligned}$$



- c) Find the equation of the tangent to the circle  $x^2 + y^2 - 8x - 14y + 40 = 0$  at the point (8,4)

$$\begin{aligned} (x-4)^2 - 16 + (y-7)^2 - 49 + 40 &= 0 \\ (x-4)^2 + (y-7)^2 &= 25 \\ \text{centre } (4, 7) & \text{ radius } 5 \\ \text{gradient radius} &= \frac{-3}{4} \\ \text{to } (8, 4) & \\ \text{gradient tangent} &= \frac{4}{3} \\ y - 4 &= \frac{4}{3}(x - 8) \\ 3y - 12 &= 4x - 32 \\ 4x - 3y &= 20 \end{aligned}$$

- d) A tangent to the circle  $x^2 + y^2 = 36$  passes through the point (10,0) and crosses the positive y-axis. What is the coordinate of the point where the tangent meets the y-axis?



$$\begin{aligned} y &= mx + c \\ 0 &= 10m + c \\ c &= -10m \\ y &= m(x - 10) \\ y^2 &= m^2(x^2 - 20x + 100) \\ c &= \frac{15}{2} \\ x^2 + m^2(x^2 - 20x + 100) &= 36 \\ x^2(1+m^2) - 20m^2x + (100m^2 - 36) &= 0 \\ \Delta &= 0 \quad 400m^4 - 4(1+m^2)(100m^2 - 36) = 0 \\ 400m^4 - 400m^2 - 400m^4 + 144 + 144m^2 &= 0 \\ -100m^2 + 36 + 36m^2 &= 0 \\ 64m^2 &= 36 \quad m^2 = \frac{9}{16} \quad m = \pm \frac{3}{4} \\ m &= -\frac{3}{4} \quad y = -\frac{3}{4}x + \frac{15}{2} \quad (0, \frac{15}{2}) \end{aligned}$$

- e) Find the radius of the circle with equation  $2x^2 + 2y^2 + 12x - 4y + 13 = 0$

$$\begin{aligned} x^2 + y^2 + 6x - 2y + \frac{13}{2} &= 0 \\ (x+3)^2 - 9 + (y-1)^2 - 1 + \frac{13}{2} &= 0 \\ (x+3)^2 + (y-1)^2 &= \frac{7}{2} \\ \text{radius} &= \sqrt{\frac{7}{2}} \end{aligned}$$

- f) A circle has equation  $x^2 + y^2 - 10x - 12y + 56 = 0$  and  $C$  is the centre of the circle.  
The tangent to the circle at  $A(6,4)$  meets the  $y$ -axis at  $B$ . Find the area of triangle  $ABC$ .

$$(x-5)^2 - 25 + (y-6)^2 - 36 + 56 = 0$$

$$(x-5)^2 + (y-6)^2 = 5 \quad (5,6)$$

grad AC = -2  
grad AB =  $\frac{1}{2}$   
 $y - 4 = \frac{1}{2}(x - 6)$   
 $y = \frac{1}{2}x + 1$

$$|AB| = \sqrt{6^2 + 3^2} = \sqrt{45}$$

$$|AC| = \sqrt{5}$$

$$\text{Area } ABC = \frac{1}{2} \times \sqrt{45} \times \sqrt{5}$$

$$= \frac{15}{2}$$

- g) A circle has centre  $(8,k)$  where  $k$  is a constant.  
The straight line with equation  $y = 3x - 12$  is tangent to the circle at  $(5,3)$ .  
Find the equation of the circle.

grad radius =  $\frac{k-3}{3} = -\frac{1}{3}$      $k=2$     Centre  $(8,2)$

$$r^2 = 3^2 + 1^2 = 10$$

$$(x-8)^2 + (y-2)^2 = 10$$

- h) A circle has centre  $(5,6)$ .  
The straight line which passes through  $(1,8)$  and  $(10,11)$  is a tangent to the circle.  
Find the radius of the circle.

gradient of tangent =  $\frac{3}{9} = \frac{1}{3}$      $y - 8 = \frac{1}{3}(x - 1)$   
 $y = \frac{1}{3}x + \frac{23}{3}$

gradient of radius = -3  
 $y - 6 = -3(x - 5)$   
 $y = -3x + 21$

$$-3x + 21 = \frac{1}{3}x + \frac{23}{3}$$

$$-9x + 63 = x + 23$$

$$10x = 40 \quad x = 4$$

$$y = 9$$

$$r = \sqrt{1^2 + 3^2}$$

$$= \sqrt{10}$$

- i) A circle has equation  $x^2 + y^2 + 2x - 4y + 1 = 0$ .  
The straight line with equation  $y = mx$  is a tangent to the circle.  
Find the difference in the possible values of  $m$ .

$$(x+1)^2 + (y-2)^2 = 4 \quad y = mx$$

$$(x+1)^2 + (mx-2)^2 = 4$$

$$x^2 + 2x + 1 + m^2x^2 - 4mx = 0$$

$$(1+m^2)x^2 + (2-4m)x + 1 = 0$$

$$\Delta = 0 \quad (2-4m)^2 - 4(1+m^2) = 0$$


$$1 - 4m + 4m^2 - 1 - m^2 = 0$$

$$3m^2 - 4m = 0$$

$$m(3m - 4) = 0$$

$m = 0, \frac{4}{3}$

$$\frac{4}{3}$$



j) A circle has centre at the origin and radius  $R$ .  $x^2 + y^2 = R^2$   
 The circle fits wholly inside the circle with equation  $x^2 + y^2 - 10x - 24y = 231$ .  
 Find the range of possible value of  $R$ .

$$(x-5)^2 + (y-12)^2 - 25 - 144 = 231$$

$$(x-5)^2 + (y-12)^2 = 20^2$$

Distance  $O \rightarrow$  centre  $= \sqrt{5^2 + 12^2} = 13$

$$R \leq 20 - 13 \quad R \leq 7 \quad \underline{\underline{0 < R \leq 7}}$$

k) A circle is drawn inside a regular hexagon so that the circle touches each side of the hexagon.  
 What fraction of the hexagon is covered by the circle?

Circle:  $\pi r^2$

fraction:  $\frac{\pi r^2}{2\sqrt{3}r^2}$

$$\frac{\pi}{2\sqrt{3}} = \underline{\underline{\frac{\sqrt{3}\pi}{6}}}$$

Circle:  $\frac{a^2}{4} + r^2 = a^2$   
 $r^2 = \frac{3a^2}{4} \quad a^2 = \frac{4r^2}{3}$

Hexagon:  $6 \times \frac{1}{2} a^2 \frac{\sqrt{3}}{2} = 6 \times \frac{1}{2} \times \frac{4r^2}{3} \times \frac{\sqrt{3}}{2} = 2\sqrt{3}r^2$

l) Find the shortest distance between the circle  $x^2 + y^2 + 6x + 8y = 75$  and the origin.

$$(x+3)^2 + (y+4)^2 = 100$$

gradient  $= \frac{4}{3}$   
 $y = \frac{4}{3}x$

$$x^2 + \frac{16}{9}x^2 + 6x + \frac{32}{3}x = 75$$

$$\frac{25x^2}{9} + \frac{50x}{3} - 75 = 0$$

$$x^2 + 6x - 27 = 0$$

$$(x-3)(x+6) = 0$$

$$x = 3 \quad y = 4$$

(3,4)  
 Distance  $= \sqrt{3^2 + 4^2} = \underline{\underline{5}}$

m) Find the shortest distance between the line  $x + 2y = 2$  and the circle  $x^2 + y^2 - 6x - 8y + 21 = 0$

$$(x-3)^2 + (y-4)^2 = 4$$

1<sup>st</sup> gradient  $= 2$  (3,4)  
 $y - 4 = 2(x - 3)$   
 $y = 2x - 2$   
 $2x - 2 = -\frac{1}{2}x + 1$   
 $\frac{5}{2}x = 3 \quad x = \frac{6}{5} \quad y = \frac{2}{5}$

Distance line to centre  
 $= \sqrt{(\frac{18}{5})^2 + (\frac{9}{5})^2}$   
 $= \frac{9}{5} \sqrt{4+1} = \frac{9\sqrt{5}}{5}$   
 Distance line to circle  
 $= \underline{\underline{\frac{9\sqrt{5}}{5} - 2}}$

n) Find the shortest distance between the two circle with equations:


$$(x-5)^2 + (y-9)^2 = 45 \quad \text{and} \quad (x+1)^2 + (y+3)^2 = 5$$

(5, 9)  $r = 3\sqrt{5}$       (-1, -3)  $r = \sqrt{5}$

distance between centres  $= \sqrt{6^2 + 12^2} = 6\sqrt{5}$

distance between circles  
 $= 6\sqrt{5} - 3\sqrt{5} - \sqrt{5}$   
 $= \underline{\underline{2\sqrt{5}}}$

- o) The two circles with equations below have exactly one point in common. Centre  $(-1, 5)$   $r = 6$   
 Centre  $(8, -7)$  radius  $r$
- $$(x + 1)^2 + (y - 5)^2 = 36 \quad \text{and} \quad (x - 8)^2 + (y + 7)^2 = r^2$$
- Find the two possible values of  $r$




Distance between centres =  $\sqrt{9^2 + 12^2} = 15$

$$r_1 = 15 - 6 = 9$$

$$r_2 = 15 + 6 = 21$$

- p) The two circles with equations below have exactly one point in common.
- $$(x + r)^2 + (y + r)^2 = 4r^2 \quad \text{and} \quad (x - r)^2 + (y - 2)^2 = r^2$$
- Find the value of  $r$ , where  $r > 0$



Distance between centres =  $3r$

$$(r + 2)^2 + (2r)^2 = 9r^2$$

$$r^2 + 4r + 4 + 4r^2 = 9r^2$$

$$4r^2 - 4r - 4 = 0$$

$$r^2 - r - 1 = 0$$

$(-r, -r)$   $(r, 2)$



$$r = \frac{1 \pm \sqrt{1 + 4}}{2} \quad r > 0$$

$$r = \frac{1 + \sqrt{5}}{2}$$

- q) Circle  $C_1$  has equation  $(x + 2)^2 + (y - 2)^2 = 7$   $(-2, 2)$   $r = \sqrt{7}$   
 Circle  $C_2$  has equation  $(x - 6)^2 + (y - 2)^2 = 7$   $(6, 2)$


The straight line  $L$  is a tangent to both circles and has a positive gradient.  
 The angle between  $L$  and the x-axis is  $\theta$ . Find  $\cos \theta$

Distance between centres = 8  
 $OM = 4$   
 $k^2 = 16 - 7 = 9$   
 $k = 3$   
 $\cos \theta = \frac{3}{4}$



- r) Circle  $C_1$  has equation  $x^2 + y^2 - 10x - 10y + 41 = 0$   
 Circle  $C_2$  has centre  $(k, 5)$  and touches both  $C_1$  and the y-axis  
 Find the difference between the two possible values of  $k$ .

$C_1$   $(x - 5)^2 + (y - 5)^2 = 9$   $r = 3$



small circle  $d = 2, r = 1$   $k = 1$   
 large circle  $d = 8, r = 4$   $k = 4$

Difference = 3